

AERO® line of Insecticide Products contains a full range of ready-to-use residual and space spray products for all indoor and outdoor pest control needs.



aero

Assault II

- **INTENDED FOR COMMERCIAL USE ONLY • A READY-TO-USE STABLE EMULSION INSECTICIDE**
- **FOR USE OUTDOORS AND NON-FOOD AREAS OF FOOD/FEED PROCESSING ESTABLISHMENTS, KENNELS, COMMERCIAL BUILDINGS, WAREHOUSES, THEATERS, OFFICE BUILDINGS, SCHOOLS, MOTELS, AND HOTELS.**
- **FOR CONTROL OF FLEAS, BROWN DOG TICKS AND LICE ON PREMISES**
- **FOR CONTROL OF TICKS WHICH MAY CAUSE LYME DISEASE ON PREMISES**
- **FOR CONTROL OF PESTS ON LIVESTOCK AND LIVESTOCK PREMISES**
- **PROVIDES 60-DAY (Two Months) CONTROL OF COCKROACHES**
- **PROVIDES 60-DAY (Two Months) CONTROL OF FLEAS**
- **PROVIDES 60-DAY (Two Months) CONTROL OF STORED PRODUCT PESTS**

INDOOR APPLICATION

Nonfood/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended. Do not apply to classrooms while in use.

SURFACE SPRAYING: TO CONTROL COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDES, CENTIPEDES, SOWBUGS, PILLBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Allow articles to dry thoroughly before use.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Allow to dry thoroughly before allowing dogs to contact treated areas. IF REINFESTATION OCCURS REPEAT TREATMENT BUT DO NOT REPEAT MORE FREQUENTLY THAN EVERY 14 DAYS.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of *House Flies*, *Fruit Flies*, *Gnats*, *Mosquitoes*, *Skipper Flies*, *Wasps*, *Hornets*, *Bees*, *Black Flies*, *Angoumois Grain Moths*, *Mediterranean Flour Moths*, *Indian Meal Moths* and *Tobacco Moths*: Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for 2 hours. Ventilate before reoccupying.

OUTDOOR APPLICATION

TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT): Use as a residual treatment to control Ticks including the *Deer Tick (Ixodes dammini)*, *Lone Star Tick (Amblyomma americanum)*, *Brown Dog Tick (Rhipicephalus sanguineus)*, *American Dog Tick (Dermacentor variabilis)* and *Western Black-legged Tick (Ixodes pacificus)*. For each area listed below, apply until surface is wet (approximately 1 gal/750-1000 ft²). Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces where these pests may enter. Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control ticks occurring there. To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

Distributed by: AERO Chemical Company
PO Box 16247, Atlanta, GA 30321-0247
AERO is a registered trademark of ABC Compounding Co., Inc.

HIGHLIGHTS

- Ready to use
- Residual: up to 60 days
- Water-based
- Effective against fleas and ticks
- Effective against ants and cockroaches
- May be used through a fogger
- Ornamental plant use
- Mosquito control

TYPICAL PROPERTIES

- Specific Gravity: 1.0
- pH: 7.5 - 10
- **ACTIVE INGREDIENTS:**
Permethrin0.50%
Other Ingredients.....99.50%
- Appearance: White emulsion
- Odor: Insecticidal

SAFETY CAUTIONS

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash before reuse.

Refer to SDS and product label for additional safety information and directions for use.